

Table 18. Reported cases of HIV infection (not AIDS), by age category, transmission category, and sex, cumulative through 2004—42 areas with confidential name-based HIV infection reporting

Transmission category	Males				Females				Total			
	2004		Cumulative through 2004 ^a		2004		Cumulative through 2004 ^a		2004		Cumulative through 2004 ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Adult or adolescent												
Male-to-male sexual contact	11,601	50	76,003	48	—	—	—	—	11,601	35	76,003	34
Injection drug use	2,675	12	19,989	13	1,267	13	11,719	17	3,942	12	31,709	14
Male-to-male sexual contact and injection drug use	878	4	8,298	5	—	—	—	—	878	3	8,298	4
Hemophilia/coagulation disorder	29	0	474	0	7	0	62	0	36	0	536	0
Heterosexual contact	2,014	9	13,564	9	4,090	41	30,644	45	6,104	18	44,208	20
<i>Sex with injection drug user</i>	293	1	2,330	1	666	7	6,063	9	959	3	8,393	4
<i>Sex with bisexual male</i>	0	0	0	0	189	2	1,822	3	189	1	1,822	1
<i>Sex with person with hemophilia</i>	6	0	29	0	14	0	182	0	20	0	211	0
<i>Sex with HIV-infected transfusion recipient</i>	17	0	124	0	26	0	186	0	43	0	310	0
<i>Sex with HIV-infected person, risk factor not specified</i>	1,698	7	11,081	7	3,195	32	22,391	33	4,893	15	33,472	15
Receipt of blood transfusion, blood components, or tissue	46	0	468	0	64	1	546	1	110	0	1,014	0
Other/risk factor not reported or identified	6,015	26	38,251	24	4,446	45	24,572	36	10,461	32	62,829	28
Subtotal	23,258	100	157,047	100	9,874	100	67,543	100	33,132	100	224,597	100
Child (<13 yrs at diagnosis)												
Hemophilia/coagulation disorder	4	2	106	4	0	0	1	0	4	1	107	2
Mother with the following risk factor for, or documented, HIV infection	160	78	1,991	84	184	81	2,123	87	344	80	4,114	85
<i>Injection drug use</i>	25	12	523	22	23	10	532	22	48	11	1,055	22
<i>Sex with injection drug user</i>	10	5	200	8	15	7	200	8	25	6	400	8
<i>Sex with bisexual male</i>	5	2	30	1	2	1	21	1	7	2	51	1
<i>Sex with person with hemophilia</i>	1	0	3	0	0	0	7	0	1	0	10	0
<i>Sex with HIV-infected transfusion recipient</i>	1	0	6	0	0	0	4	0	1	0	10	0
<i>Sex with HIV-infected person, risk factor not specified</i>	45	22	457	19	44	19	524	22	89	21	981	20
<i>Receipt of blood transfusion, blood components, or tissue</i>	0	0	17	1	1	0	17	1	1	0	34	1
<i>Has HIV infection, risk factor not specified</i>	73	36	755	32	99	44	818	34	172	40	1,573	33
Receipt of blood transfusion, blood components, or tissue	2	1	25	1	1	0	23	1	3	1	48	1
Other/risk factor not reported or identified	39	19	262	11	41	18	283	12	80	19	545	11
Subtotal	205	100	2,384	100	226	100	2,430	100	431	100	4,814	100
Total	23,463	100	159,431	100	10,100	100	69,973	100	33,563	100	229,411	100

Note. Includes only persons with HIV infection that has not progressed to AIDS.

Since 2004, the following 42 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Georgia, Idaho, Indiana, Iowa, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Puerto Rico, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, American Samoa, Guam, Northern Mariana Islands, and the U.S. Virgin Islands.

Connecticut has confidential name-based HIV infection reporting only for pediatric cases.

Florida (since July 1997) has had confidential name-based HIV infection reporting only for new diagnoses.

Pennsylvania (October 2002) implemented confidential name-based HIV infection reporting only in areas outside the city of Philadelphia.

Texas (February 1994 through December 1998) reported only pediatric HIV infection cases.

^a Includes persons with a diagnosis of HIV infection (not AIDS), reported from the beginning of the epidemic through December 2004. Cumulative total includes 7 persons of unknown sex.